

# Formations of endogenous ore deposits and mineralization in Vietnam

Nguyen Van Nhan

*College of Science, VNU*

Received 14 January 2008; received in revised form 05 March 2008

**Abstract.** Formations of endogenous ore deposits and mineralization are widely distributed in Vietnam. Detailed studies allow for distinguishing 30 ore formations representing 9 groups of mineral raw - materials: Cr, Ti, Fe, Sn - W, Mo, Pb - Zn, Au, Sb - Hg. Three of presented formations are of magmatic affiliation, one is pegmatitic, three others belong to skarn mineralization and the remaining ones are of hydrothermal - volcanic and controversial origin. Three metallogenic epochs can be distinguished: Proterozoic, Palaeozoic and Mesozoic - Cainozoic. Taking into consideration the recent structural data and economic importance along with the geochemical units can be distinguished in Vietnam: 1. Sialic - mafic Viet Bac; 2. Mafic Tay Bac; 3. Sialic Truong Son; 4. Mafic - metamorphic Kontum; 5. Sialic - alkaline Dalat - Nam Bo.

**Keywords:** Endogenous ore deposit; Mineralization.

Vietnam and other countries of SE Asia cover an outstanding area of the crossing of two structural and metallogenic belts: Pacific and Mediterranean. Simultaneously, this area is an active continental margin. Abundant and variable ore mineralization is known with the famous W - Sn belt and deposits of Cr, Cu, Zn - Pb, Au, Sb - Hg, Bi, Mo and REE.

Geological history of the investigated area is influenced by the development of the two large adjacent structures: Southern China platform and Indo - China fold belt. Typical feature of Vietnam area is the Mesozoic and Cainozoic activation structures. Numerous intrusions ranging in age from Proterozoic to Cainozoic were formed in tectono - magmatic

cycles operating within the fold belts and activated areas.

Detailed studies allow for distinguishing 30 ore formations representing 9 groups of mineral raw - materials: Cr, Ti, Fe (Fig. 1), Sn - W (Fig. 2), Mo, Zn - Pb, Au, Sb - Hg (Fig. 3). Three of presented formations are of magmatic affiliation, one is pegmatitic, three others belong to skarn mineralization and the remaining ones are of hydrothermal - volcanic and controversial origin.

Three metallogenic epochs can be distinguished: Proterozoic, Palaeozoic and Mesozoic - Cainozoic. Archean endogenous mineralization is poorly known but others contain numerous, genetically diversified magmatic and post - magmatic deposits.

\* Tel.: 84-984973786.




Fig. 1. Scheme of Cr, Ti, Fe ore formations in Vietnam

**Metallogenic units:** I. Viet Bac; II. Tay Bac; III. Truong Son; IV. Kon Tum; V. Da Lat - Nam Bo; VI. The Cr ore deposits (1. Nui Nua, 2. Hon Vang); VII. The Ti ore deposits (3. Nui Chua, 4. Tam ki, 5. Xa Hieu); VIII. The Fe ore deposits (6. Cao Bang, 7. Ha Tinh, 8. Trai Cau, 9. Lang Khuan, 10. Linh Nham).

During Proterozoic the area of Vietnam was a proto - geosynclinal - proto continental area marked by basaltoid - granitoid magmatism. In the initial geosynclinal stage sedimentary - volcanic pile of considerable thickness was deposited. Basic volcanic and meta - gabbros (ophiolite type) contain chalcopyrite - magnetite - ortite formation (s.c., Cu - pyrites) within metasomatic rocks of Proterozoic basement. Pyrite and polymetallic Cu - pyrite formations genetically

connected with basic metavolcanics (greenschists) also belong to this stage.

Palaeozoic metallogenic epoch embraces several ore - formations. Few ultrabasic - basic intrusions contain Cu - Ni and Ti - magnetite mineralization. Pegmatites related to Late - Palaeozoic granite intrusions host cassiterite formation with accompanying Nb and Ta. Galena - sphalerite formation with Pb - Sb sulphosalts is connected with granite - grano - syenite intrusions and associated liparite - trachites.




Fig. 2. Scheme of Sn - W ore formations in Vietnam.

**Metallogenic units:** I. Viet Bac; II. Tay Bac; III. Truong Son; IV. Kon Tum; V. Da Lat - Nam Bo; VI. The Sn - W ore deposits (39. Tinh Tuc, 40. Tam Dao, 41. Quy Hop, 42. Kim Cuong, 43. Ban Chieng, 44. Phu Loi).

Post - Permian rejuvenation of tectonic and magmatic activity in the whole area of Vietnam resulted in numerous valuable ore and other raw - material deposits. Two metallogenic epochs can be distinguished: Cimmerian and Alpine.

Cimmerian epoch includes deposits and mineralization of Fe, Cr, Cu, Ni, Pb - Zn, Mo, Sn, W, Sb - Hg, Au - Ag. In the Black River depression Permo - Triassic basic volcanics are common with the related formations of pyrites, Cu - pyrites, chalcopyrite - quartz and native copper. Magnetite - hematite hydrothermal - metasomatic formation can also be connected with this basic volcanism. Chromite formation occurs in Permo - Triassic apo - harzburgite intrusions located along the deep fault structure of Ma River. Ilmenite - Ti - magnetite mineralization is known from the widespread Upper Triassic basic intrusions, whereas Cu - Ni formation is hosted in Permo - Triassic serpentized dunites. Granite - granodiorite complexes (Permo - Triassic) provide numerous other formations: sphalerite - galena in carbonate rocks, sphalerite - galena - pyrite - arsenopyrite, and galena - quartz in granites, magnetite in skarns and syderite. Granodiorite - mozonite intrusions (100 - 190 my) contain molybdenite - chalcopyrite formation. Sub - volcanic intrusions accompanied by acid - alkaline volcanics (Jurassic - Cretaceous) are connected with galena - sphalerite formation with Pb - Sb sulphosalts.




Fig. 3. Scheme of Cu, Pb - Zn, Au, Sb - Hg, Mo ore formations in Vietnam.

**Metallogenic units:** I. Viet Bac; II. Tay Bac; III. Truong Son; IV. Kon Tum; V. Da Lat - Nam Bo; VI. The Cu ore deposits (11. Sin Quyen, 12. Son La, 13. Cao Bang, 14. Tra Lam, 15. Van Sai, 16. Luong Son, 17. Nui Nua, 18. Bac Giang); VII. The Pb - Zn ore deposits (19. Ngan Son, 20. Cho Dien, 21. Lang Hit, 22. Quan Son, 23. Mi Duc, 24. Tu Le, 25. Na Son, 26. Duc Bo, 27. Phu Loi); VIII. The Au ore deposits (28. Bong Mieu, 29. Pac Lan); IX. The Sb - Hg ore deposits (30. Chiem Hoa, 31. Ba Thuoc, 32. Dien Bien, 33. Bac Thai, 34. Yen Ve); X. The Mo ore deposits (35. Sapa, 36. Nui Sam, 37. Krongfa, 38. Nui Dat).

Alpine metallogenic epoch is characterized by ore deposits of Sn, Mo, Zn - Pb, Ag - Au, Sb - Bi, U and REE with the leading deposits of Sn, W, Mo and REE. Sphalerite - galena - cassiterite formation rich in Ag along with Au - quartz - sulphide formation are connected with Upper - Cretaceous granites and leucogranites. Au - Sb formation is spatially related to the Sn and Zn - Pb - Ag ores. Tin deposits belonging to cassiterite - wolframite - quartz and cassiterite

- silicate - sulphide formations are hosted in Upper Cretaceous two - mica granites. In Southern Vietnam quartz - molybdenite veins cut Cretaceous leucocratic - biotitic granite massifs. Palaeogene granite - granosyenite intrusions provide Sn and Zn - Pb ores belonging to cassiterite - silicate - sulphide, cassiterite and galena - sphalerite formations in skarns, molybdenite - quartz formation and molybdenite - uraninite one.

Table 1. Metallogenic epochs in Vietnam

Epochs	Ore complex	Ore formation	Ore deposits
PR - PZ1	Relate to metasomatic rocks	- Chalcopyrite - magnetite - orthite (Cu - pyrite)	Sin Quyen
	Relate to green schist	- Pyrite - Polymetalic Cu - pyrite	Bo Xinh, Duc Phu, Duc Bo
PZ	Relate to ultramafic - mafic intrusions	- Pyrrhotite - pentlandite - chalcopyrite - Ilmenite Ti - magnetite	Tra Lam, Dac Sa Tam Ky, Xa Hieu
	Relate to granitoid	- Cassiterite in pegmatite	Kim Cuong
	Relate to alkaline acid extrusion	- Galena - sphalerite formation with Pb - Sb sulphosalts	Na Son
MZ - KZ  Cimmerian	Relate to basic extrusion	- Pyrite - Pyrite - Zn - Pb - Cu - Chalcopyrit - quartz - Native copper - Magnetite - hematite	Song Da Van Sai, Quy Huong Ban Mua Trai Cau
	Relate to apo - harzburgite	- Chromite	Co Dinh
	Relate to dunite, gabbro, peridotite	- Pyrrhotite - pentlandite - chalcopyrite	Ban Xang, Cao Bang
	Relate to gabbro	- Ilmenite Ti - magnetite	Cay Cham
	Relate to granite, grano diorit	- Dien Bien - Phiabroc  - Van Canh - Phu Trai	Dien Bien Cho Dien  Thai Nguyen Ha Tinh, Na Rua Ban Phang Tien Thuan, An Khe
	Unknown relate to magma	- Bornite - chalcocite - Antimonite - quartz	Bac Giang, Mau Son Quang Ninh
	Relate to granodiorite - mozonite	- Molybdenite - chalcopyrite - Galena - quartz in granite	Krongfa Gia Bac

Epochs		Ore complex	Ore formation	Ore deposits	
		Relate to subpluton and alkaline acid extrusion	- Galena - sphalerite formation with Pb - Sb sulphosalts	Tu Le	
		Relate to granite - granosyenite	- Cassiterite - wolframite - quartz - Molybdenite - wolframite	Mepu Nui Dat	
		Relate to basanite andesite	Pyrite with Au, Ag	Thu Duc	
MZ - KZ	Alpine	Relate to granite leucogranite	- Piaoak - Deo Ca	- Sphalerite - galena - cassiterite formation rich in Ag - Au - quartz - sulphide - Cassiterite - wolframite - quartz - Cassiterite - silicate - sulphide - Au - antimonite - Molybdenite - quartz	Ngan Son Pac Lan Tinh Tuc Tam Dao Chiem Hoa Nui Sam
				- Cassiterite - wolframite - quartz - Cassiterite - silicate - sulphide - Cassiterite in skarn	Bu Me Quy Hop Phu Loi
		Relate to granite, granosyenite	Song Chu, Ban Chieng Yeyensun	- Galena - sphalerite in skarn - Molybdenite - quartz - Molybdenite - uraninite - Chalcopyrite with U, REE	Ke Tang Sapa Vi Kem Sin Quyен

Taking into consideration recent structural data and economic importance along with the geochemical character of mineralizations, five large mineralogical units can be distinguished in Vietnam as follows:

1. Sialic - mafic Viet Bac;
2. Mafic Tay Bac;
3. Sialic Truong Son;
4. Mafic - metamorphic Kon Tum;
5. Sialic - alkaline Da Lat - Nam Bo.

The two first units belong to Vietnam - Chon fold belt whereas the remaining represents Indo - China fold belt. Apart from that, two separate zones were distinguished: Muong Te and Ha Tien which are the parts of Thailand - Malay geosynclinal - fold belt. Each unit reveal typical metalogenic features controlled by tectonic - structural development:

1. Viet Bac: Sn - W, Zn - Pb, Au, Sb - Hg;
2. Tay Bac: Cu, Cu - Ni, pyrites, Cr, Mo, REE, Zn - Pb;
3. Truong Son: Sn - W, Zn - Pb, Fe, Au, Sb - Hg;
4. Kontum: Au;
5. Dalat - Nam Bo: Mo, Bi.

Endogenous ore - forming processes are clearly confined to the magmatic and tectonic activity in the specific geological epochs. The first group embraces deposits of Cr, Ni, Co, Cu, Cu - Ni, Fe (contact - metasomatic), Ti and pyrites, all of them connected with basic/ultrabasic intrusions and located along the faults. The second group embraces polymetallic Zn - Pb, Ag, Sb, Hg, Au, Sn - W and rare - metal deposits genetically related to granites and/or granodiorites and occurring within the tectonic blocks or, commonly, at the contacts between two areas of different structural and facial characters. The crucial factor controlling the metallogeny of this territory was magmatic/tectonic activation cycle in Mesozoic and Cainozoic.

Endogenous mineralization in Vietnam bears the metallogenetic features of the Pacific belt with typical Sn - W, Mo, Zn - Pb and Au deposits. Mediterranean belt features can be observed in Tay Bac unit with the typical Cu, Cu - Ni, pyrites, Cu - polymetallic pyrite deposits.

## References

- [1] Dinh Van Dien et al., *Minerals and resources of Vietnam - general of development of Geology and Minerals of Vietnam*, Department of Geology and Minerals of Vietnam, Hanoi, 1995, 254.
- [2] Nguyen Van Nhan, *Polymetallic Ore formations in North of Vietnam*, IV, GEOSEA, Malina, Philipin, 1981.
- [3] Nguyen Van Nhan, Formations of endogenous ore deposits and mineralization in Vietnam, Scientific bulletins of Stanislaw staszic Academy of Mining and Metallurgy, Krakow, Poland, *Geology bulletin* No. 38 (1988) 134.
- [4] Nguyen Van Nhan, Formations of endogenous ore deposits in Vietnam, *Journal of Geology, series A*, No. 234 (1996) 7 (in Vietnamese).