

Cenozoic magmatism of Eastern Sea (South China Sea)

Phan Truong Thi¹, Vo Viet Van^{2,*}

¹ College of Science, VNU Hanoi

² University of Technology, VNU Ho Chi Minh City

Received 07 January 2008; received in revised form 12 March 2008

Abstract. In Eastern Sea (South China Sea), Jurassic - Cretaceous magmatic intrusives are discovered, they were cutting the Pre-Cenozoic sedimentaries. The Cenozoic volcanic basalts are strongly distributed together with Cenozoic sedimentaries. The paper concerns only the volcanic activities with description on their geology, chemistry, geochemistry, isotopic ages and their forming mechanism. In fact, the main topic of the paper is dealt with the forming mechanism of basalt in the dynamics of the opening of Eastern Sea in particular, and of South China Sea in general.

Keywords: Cenozoic; Magmatism; Eastern Sea.

1. Generality on cenozoic geology of Eastern Sea

The geomorphology and general distribution picture of Eastern Sea crust types are presented in Fig. 1. The given data are received from the research of satellite images and of geophysics measurements. The crusts are covered by Cenozoic sedimentaries [2, 3]. Their geomagnetic ages are also represented in Fig. 2 [2, 3]. The geomagnetic anomalies lines N.5 (16.5 Ma) and N.13 (32 Ma) are remarked, they also are recognized as the timing of the opening and closing of Eastern Sea in his geodynamics respectively. The follows are Late Miocene and Pliocene sedimentaries, covered all the territory of the South China Sea. These results are obtained




Fig. 1. The geomorphology and structure of crust types of South China Sea.

* Corresponding author. Tel.: 84-913535203
E-mail: banganus@yahoo.com

by scientific cooperation between Hanoi University and Paris VI University [8, 9, 12].

2. Petrography and petrochemistry of Eastern Sea basalts

According to the research data of seismic measurements on the Eastern Sea, the volcanic basalts mainly cut out the sedimentaries of Oligocen - Pleistocen. Geomorphological study of the bottom of the sea also makes clear the strong relief character by a lot of underground volcanic mountains.

After the study of volcanic isotopic ages by method K-Ar and by track fission of zircon grains, the ages of basalts varied mainly from 13 Ma to 1924 year. It is clear in Table 1 that the age of basalt on the Re Island is 13 Ma, meanwhile the age of the basalt in Con Co Island is only 3,200 years. In the 1924 year, there was a volcanic activity near Nha Trang City.

From the petrochemical data (Tables 1

and 2) it is clear that basalts of Eastern Sea are belonged to two rock types: alkaline and tholeiitic series. The first one is characterized by the Ne-normative mineral, and the second one - by the positive Q- normative and absent of Ne-normative mineral.

Geologically, the basalt bodies mainly cut up the sedimentaries of Oligocene and Middle Miocene ages, they have isotopic ages varied from 13 Ma up to day (1924). It means that they are more young than the closing timing of Eastern Sea (16.5 Ma). But after the drill core samples of Cuu Long and Nam Con Son basins, the intercalation of basalt layers among the sedimentaries also being seen (Fig. 3 [15]). Although that, mainly the volcanic activity in Eastern Sea was strongly developed after the closing time of the spreading process of Eastern Sea.




Fig. 2. Schema distribution of geomagnetic anomaly lines [2, 3].

Table 1. The chemistry compositions, isotopic ages of basalt formations of the Eastern Sea [13]

Sites	XL II	XL I	1-Cendre	1-Cendre	Kawit-Is	Re- Is 1	Re-Is-2	QN	Con Co Is	Khe Sanh	DBP
Rock type	BA	AB	AB	OT	OT	OT	OT	AB	OT	OT	OT
SiO ₂	43.26	45.01	49.5	50.4	49.44	49.87	52.9	44.77	52.67	48	55.26
TiO ₂	2.76	2.03	2.34	1.9	2.67	2.2	1.57	2.85	1.78	2.68	2.04
Al ₂ O ₃	12.28	13.84	13.51	13.91	13.87	15.32	15.01	14.17	16.5	14.61	16
FeO*	12.15	12.78	11.67	11.52	11.42	10.93	0.43	11.51	10.22	10.83	9.66
MnO	0.2	0.15	0.16	0.15	0.16	0.14	0.14	0.22	0.12	0.15	8.16
MgO	12.96	9.22	7.86	8.18	8.5	7.1	7.45	11.21	4.17	8.62	5.36
CaO	10.99	9.86	9	9.13	9.13	8.18	8.93	10.12	5.67	9.15	6.15
Na ₂ O	2.87	2.76	3.27	3.13	3.13	3.11	3.28	3.22	4.92	3.09	3
K ₂ O	1.32	2.48	2.27	7.83	1.83	2.1	0.92	0.94	3.22	1.54	2.04
P ₂ O ₅	0.94	0.56	0.42	0.38	0.38	0.57	0.29	1	0.72	0.54	0.32
K ₂ O/Na ₂ O	0.46	0.81	0.61	0.58	0.53	0.704	0.28	0.201	0.65	0.458	0.68
Zr/Y	10.9	10	7.1			7.8	5.3	11	11	9.2	9.1
Zr/Nb	236.6	251.8	262.3			164	100	263.2	4.06	200	5.98
Rb/Sr	0.17	0.08	0.06			0.03	0.05	0.07	0.08	0.01	0.22
Ba/Zr	3.64	3.29	2.3			3.25	2.24	3.9	3.35	2.93	1.75
n	2	3	6	8	8	4	3	2	1	1	1
Age(Ma)	0.4-1.1	0.4-1.1	1.27 - 0	7	7	0.4-1.2	12	7.1	0.35	1	1

* Sites : XL: Xuan Loc; QN: Quang Ngai; DBP : Dien Bien Phu.

Types: BA: bazanite; AB: Alkaline Basalt; OT: Tholeite basalt.




Fig. 3. Seismic section of Cuu Long basin reflected the "stratified layers" of volcanic bodies (black) in the Middle Miocene sediments [15].

Table 2. Normative composition of some basalts of Eastern Sea

	1	2	3	4	5	6	7
SiO ₂	51.11	50.00	46.95	52.91	49.71	48.9	46.73
TiO ₂	1.73	2.43	2.36	1.40	2.10	3.74	3.40
Al ₂ O ₃	16.01	13.98	15.87	15.10	16.22	16.23	16.70
Fe ₂ O ₃	-	-	-	-	-	-	-
FeO	9.92	10.78	11.09	8.21	10.80	10.15	10.11
MnO	0.12	0.13	0.15	0.14	0.11	0.14	0.19
MgO	4.05	5.60	7.00	6.97	6.03	3.06	6.12
CaO	5.50	8.42	7.05	9.00	10.86	9.18	9.52
Na ₂ O	4.77	3.38	2.64	3.12	3.08	3.57	3.28
K ₂ O	3.12	2.01	2.66	0.72	0.51	2.11	2.19
P ₂ O ₅	0.70	0.75	0.61	0.24	0.30	1.11	0.62
Total	97.03	97.48	96.38	97.81	99.73	98.21	98.9
Q				0.97			-
C				-			-
Or	18.44	11.88	15.72	4.26	3.02	12.48	12.95
Ab	31.82	28.6	21.89	26.40	26.06	30.21	18.07
An	13.06	17.04	23.60	25.07	28.94	22.03	24.38
Lc	-	-	-	-	-	-	-
Ne	4.63	-	0.24	-	-	-	5.25
Di	8.10	16.49	6.11	14.76	19.12	14.18	15.70
Wo	-	-	-	-	-	-	-
Hy	-	-	-	23.14	8.45	1.89	-
Ol	16.07	13.14	22.93	-	7.87	6.34	13.18
Mt	-	-	-	-	1.74	1.64	1.62
Il	3.29	4.62	4.48	2.66	3.99	7.10	6.46
Ap	1.62	1.74	1.41	0.56	0.65	2.42	1.35
Cc							

Table 3. The chemical compositions of stratified layers of basalts among the sedimentaries of Cuu Long Basin [15]

	1	2	3	4	5	6	7	8	9	10
SiO ₂	45.18	50.38	47.26	47.67	49.34	47.47	49.00	47.39	50.16	48.56
TiO ₂	1.45	0.95	1.68	1.67	1.88	1.26	1.32	1.19	1.32	1.33
Al ₂ O ₃	17.67	17.70	15.38	17.42	16.59	16.92	15.90	17.16	15.68	16.44
Fe ₂ O ₃	2.28	3.04	4.08	3.53	9.76	3.14	2.18	10.94	5.18	8.47
FeO	6.27	4.30	8.08	6.36	-	8.15	6.27	-	4.92	-
MnO	0.14	0.11	-	0.14	0.30	0.16	0.10	0.17	0.12	0.11
MgO	6.85	5.63	6.77	6.67	6.45	9.00	5.87	8.87	7.38	5.84
CaO	6.03	7.81	4.78	3.67	2.69	2.11	5.86	2.09	9.61	5.90
Na ₂ O	4.75	3.09	3.02	5.29	0.61	2.45	5.22	2.45	3.34	5.22
K ₂ O	1.38	1.42	2.10	0.66	7.84	4.06	0.79	4.41	0.10	0.82
P ₂ O ₅	0.32	0.39	0.11	0.29	0.27	0.16	0.21	0.16	0.16	0.24
S	0.10	0.10	-	0.10	-	0.10	0.10	-	0.10	0.10
LOI	6.90	4.68	6.23	6.12	3.77	4.67	6.57	4.67	1.47	6.57
Total	99.32	99.60	99.49	99.49	99.50	99.41	99.39	99.50	99.54	99.50

From Table 3, the total amount of (Na₂O + K₂O) is more than 5-6%, what confirms that the alkalinity of rocks is high and it is belonging to the alkaline basalt type.




Fig. 4. CIPW normative variation for Eastern Sea basalts in comparison with on-land volcanic activity. Remark that the basalts is from Island as Re Island, Ile des Cendres [13].

3. Discussion about the forming mechanism

For discussion about the forming mechanism of basalt volcanic activity of Eastern Sea, it is necessary to remark the two following data:

1) By calculation of formation pressure of basalts (O' Hara diagram; by barometer), the alkaline basalts were formed at the depth of more than 90 km (pressure more than 30 kbars); meanwhile, the tholeiitic basalts were formed at the depth of about 18-30 km (pressure of about 6-10 kbars) (refer to Table 1).




Fig. 5. Spider diagram for comparing basalts from the island and on-land.

Table 4. Isotopic compositions of basalts of South China Sea [10].

Samples	Sr 87/86	Nd 143/144	ε Nd	Pb 206/204	Pb 207/204	Pb 208/204
D8-2	0.703594	0.512929	5.68	18.704	15.609	38.325
D8-4	0.703561	0.512916	5.42	18.600	15.632	38.848
D9-3	0.704433	0.512922	5.54	18.667	15.535	38.677
D9-2	0.703976	0.512813	3.41	18.954	15.588	38.991
D10	0.704007	0.512805	3.26	18.875	15.593	38.931
O23-40	0.703814	0.512952	6.13	18.601	15.557	38.629
O23-37	0.703991	0.512898	5.07	18.543	15.606	38.598
23-37-7	0.703936	0.512894	4.99	18.481	15.567	38.618
23-35-6	0.704355	0.512913	5.36	18.411	15.575	38.554
75-31	0.703689	0.513035	7.72	18.521	15.520	38.402
M32283	0.704222	0.513184	10.65	17.864	15.447	37.605
Pa1-5	0.704453	0.513129	9.58	17.886	15.449	37.648

2) The isotopic study (Table 4) all of the isotopic ratios (Sr-87/86; Nd 143/144; Pb 206/204; Pb 207/204; Pb 208/204) confirms that the alkaline basalt belonged to the partial

melting of type PREMA (PREvalent MANTle composition), it means that they are not belonged to the Depleted (DM) or Enriched Mantle (EM) (Fig. 6).




Fig. 6. Corellation diagram of isotopic ratios Nd 143/144 and Sr 86/87.

There are two problems to be discussed:

1. Why the volcanic activity has strongly developed only after the time of closing spreading process ?
2. Why there are two basalt types - alkaline and tholeiitic - of different formation depths (from 90 km to 20 km) during a single geodynamic process ?

Historically, there were a lot of hypotheses explaining geodynamics of the formation of South China Sea [2, 3, 11, 12, 14, 17-22].

A new hypothesis is proposed by the authors as follows:

The main cause of the formation of thermal dome for rising up mantle together with partial melting is the appearance of big mantle plume near the margin of Asian continent at that time.

At first, the mantle plume during its rising

up gives rise the subduction to the Borneo site (Palawan subduction), then, the subduction was extending the continental crust at the margin of Asia continent. In the depth of the Mantle, the partial melting has started but the magma liquidus was else lay with the batch melting in the depth (after the form of spider diagram). In sequences, alkaline basalt magma was formed.

During the rising up of the mantle plume, magma liquidus reservoirs from the depth were moving to the surface but not extruding on the Earth surface. At the small depth near the surface, magma liquidus reservoir became tholeiitic magma feature with positive Sr anomaly (Fig. 5). But the partial melting mechanism else remained the batch melting character (after the form of spider diagram).

When the opening process of Eastern Sea was stoped, but the magma still remains in the depth of about 20 km but not extrudes on the surface.

Additionally, under influence of the collision between the two continents India and Eurasia, a big strike slip of north-south direction (extended from Red River fault) was appeared, which moved the opening axis of Eastern Sea to the south. In consequenses, the oceanic crust was outcropped in the form of a triangle. So, the continental crust even has became thinked but covers all the surface of the south part of Eastern Sea. In these conditions, the magma liquidus reservoirs were lain in the depth. Only after closing of the rifting process, the new fault - normal or thrust kinds (from Oligocen) were opening the canals for strong volcanic activity. So, the volcanic rock types of South China Sea are very different from the magma activity kind of the all marginal seas.




Fig. 7. Some images of volcanic activity on Con Co Island (Quang Tri Province). Photographer: Phan Truong Thi. 11-1. Basalt flow; 11-2. Basalt layer with a lot of big bubbles; 11-3. Neogene sedimentary; 11-4. Image of a volcanic apparatus filled by sea water; 11-5. From the volcanic apparatus there is a basalt flow; 11-6. Neogene sedimentary (light brown color) was lain under a basalt flow (grey color); 11-7. A volcanic bomb among the sedimentary; 11-8. Sedimentary with horizontal layers; 11-9. The fault cutted sedimentaries; 11-10. The fault cutted sedimentaries.

Acknowledgements

The authors are grateful to Dr. Phan Truong Giang (Institute of Gas and Oil of Vietnam) for treatments of seismic data, Drs. Ngo Xuan Vinh, Le Van Truong and Vu Trong Hai for supplying the data of basalts in Cuu Long Basin, Dr. Nguyen Hoang and Prof. Martin Flower for discussion about the formation mechanism of basalts in South China Sea.

References

- [1] *Atlas of Geology and Geophysics of South China Sea (1/2.000.000)*, Map Publishing House, Guangdong Province (in Chinese), 1987.
- [2] A. Briaais, P. Patriat, P. Tapponier, Updated interpretation of magnetic anomalies and Seafloor spreading stages in the South China Sea. Implication for the Tertiary Tectonics of Southeast Asia, *J. Geophys. Res* 98 (1993) 6299.
- [3] A. Briaais, P. Tapponier, G. Pauttot, Constraints of Sea Beam data on crustal fabrics and seafloor spreading in the South China Sea, *Earth Plan. Sci. Letter* 95 (1989) 307.
- [4] Department of Geology and Tectonics of South China Sea, Institute of Oceanology, Academia Sinica, *South China Sea Geological Tectonics and Continental Margin Spreading*, Science Publishing House, Beijing, 1988, 376.
- [5] Gwang H. Lee, Joel S. Watkins, Seismic Sequence Stratigraphy and Hydrocarbon Potential of the Phu Khanh Basin, Offshore Central Vietnam, South China Sea, *AAPG Bull* 82 (1998) 1711-1735.
- [6] Gwang H. Lee, Keumsuk Lee, Joel S. Watkins, Geologic evolution of the Cuu Long and Nam Con Son basins, offshore southern Vietnam, South China Sea, *The Amer. Assoc. of Petrol, Geologists* 85, No. 6 (2001) 1055.
- [7] R.J.W. Gower, Early Tertiary plate reconstructions for the South China Sea region: constrains from Northwest Borneo, *Journal of Asian Earth Sciences* 4 (1990) 29.
- [8] P. Huchon, X. Le Pichon, P.T. Thi, L'équipe scientifique embarquée Campagne PONAGA, *Geochronique* 48 (1993) 9.
- [9] P. Huchon, X. Le Pichon, C. Rangin, Indochina peninsula and the collision of India and Eurasia, *Geology* 22 (1994) 27.
- [10] Kan Tu, M.F.J. Flower et al., Magmatism in the South China basin: I-Isotopic and trace element evidence for thermal erosion of the subcontinental lithosphere, *Earth and Planetary Science Letters* No. 2 (1989) 45.
- [11] Ke Ru, John D. Pigott, Episodic Rifting and Subsidence in the South China Sea, *The Amer. Assoc. of Petrol, Geologists Bull* No. 9 (1986) 70.
- [12] X. Le Pichon, P. Huchon, C. Rangin, O. Coulon, Formation of Indichinese continental margin and of the South China Sea: facts and questions, In: *Cenozoic evolution of the Indochina peninsula*, Hanoi & Do Son, Vietnam, 1995, 100.
- [13] Nguyen Hoang and Martin Flower, Petrogenesis of Cenozoic Basalts from Vietnam: Implication for Origin of a "Diffuse Igneous Province", *Journal of Petrology* 39 (1998) 369.
- [14] Nguyen Thi Ngoc Hai, *Structure et cinématique de l'extrémité de la Mer De L'Est et des bassins au sud Viet Nam*, These pour obtenir le titre de Docteur de L'Université Pierre et Marie Curie (Paris 6), 1997.
- [15] Ngo Xuan Vinh, Le Van Truong, Vu Trong Hai, Volcanic rocks of Cuu Long Basin and their reservoir characteristics. In: *The Vietnam Petroleum Institute - 25 years of its development*, Science and Technology Publisher, Hanoi, 2003, 194-214 (in Vietnamese).
- [16] G. Pauttot, C. Rangin, A. Briaais, P. Tapponier, P. Beuzart et al., Spreading direction in the central South China Sea, *Nature* 321 (1986) 150.
- [17] J.D. Pigott, K. Ru, Pulsed basin evolution on the northern margin of the South China Sea, In: *Sympo Geodynamic Evolution of the Eastern Eurasian Margin*, Paris, 1988, 88.
- [18] Phan Truong Thi, Phan Truong Dinh, Phan Truong Giang, About formatting mechanism of Eastern Sea and its oil and gaz basins. In: *The Vietnam Petroleum Institute - 25 years of its*

development, Science and Technology Publisher, Hanoi, 2003 (in Vietnamese).

- [19] K. Ru, The development of superimposed basins on the northern margin of the South China Sea and its tectonic significance, *Oil Gas Geol.* 9 (1998) 22 (in Chinese with English abstr.).
- [20] P. Tapponier, P. Pelzer, R. Armijo, On the mechanics of the collision between India and Asia, From M.P. Coward, A.C. Ries (ed), *Collision tectonics, Geol.Soc.Sp.Publi.* 19 (1986) 115.
- [21] B. Taylor, D. Hayes, Origin and geologic evolution of southern Asia Seas and Islands, part 2, AGU Washington, *Geophys.Monogr.* 27 (1983) 23.
- [22] J.M. Wu, Cenozoic basins of the South China Sea, *Episides* 11 (1988) 91.