


VNU Journal of Science: Mathematics - Physics





Original Article

The Rank of Matrices Over Positive Cones of Commutative Semirings

Ha Chi Cong*

University of Finance and Accountancy, 2 Le Quy Don, Tu Nghia, Quang Ngai, Vietnam

Received 12 December 2023 Revised 20 May 2024; Accepted 10 December 2024

Abstract: In this work, we introduced and studied the PC rank notion of a commutative semiring, and compared the PC rank with the nonnegative rank and the factor ranks of matrices, and gived also some sufficient conditions for these ranks to be the same.

Keywords: Ring, Commutative semiring, Positive cone, Idempotent matrix, Nonnegative matrix.

1. Introduction

In recent times, rank functions such as factor rank, non-negative rank and rank of completely positive matrices have attracted considerable attention from scientists when investigating conditions for a matrix to be factorizable into the product of matrices over a given commutative semiring [1-7]. In 1993, Cohen and Rothblum [1] proved some characteristic properties of the non-negative rank of nonnegative matrices over the semiring of real numbers \mathbb{R} , they also constructed a procedure for computing the non-negative rank of any non-negative matrix over this semiring. Moreover, according to [1, Theorem 4.1], if a non-negative matrix A has factor rank less than or equal to 2 then the factor rank and nonnegative rank of A coincide. In [2], Beasley and Laffey provided upper bounds on the nonnegative rank of nonnegative real matrices A and A^2 . In [3], Mohindru initiated research on completely positive matrices over commutative semirings. Accordingly, a non-negative matrix over a commutative semiring S is a matrix with entries in the positive sub-semiring P(S) of S, see [3, Definition 7.5.1]. According to [3, Proposition 7.5.2], the factor rank of a non-negative matrix is always less than or equal to its non-negative rank. In [4], Mohindru and Pereira provided necessary and sufficient conditions for a symmetric matrix over a regular incline to be completely positive, see

E-mail address: hachicong@tckt.edu.vn

Corresponding author.

[4, Theorem 2.2]. In [5], Shitov constructed an algorithm for computing the non-negative rank of tridiagonal matrices over specific commutative semirings [5, Theorem 12]. In [6], Bukovšek and Šmigoc presented some fundamental properties of symmetric non-negative matrices over the semiring of real numbers $\mathbb R$ and considered symmetric non-negative matrices trifactorization. In [7], Tung and Hoang introduced the asymptotic spectrum of non-negative matrices and some characteristic properties regarding the asymptotic non-negative rank of matrices with entries in the semiring of non-negative real numbers $\mathbb R^+$.

Regarding the rank of matrices over the positive cones of a given semiring, according to [8], every positive cone (ordering) P of a commutative semiring S always contains the positive subsemiring P(S) of S. Therefore, every matrix A with entries in P(S) can be factored into the product of two matrices over semirings P(S), P and S. For example, if matrix A has m rows and n columns then it can be factored as $A = I_m A = AI_n$, where A, I_m, I_n are matrices with entries in $P(S) \subset P \subset S$. Let k be the smallest non-negative integer such that there exist matrices B and C with entries in the positive cone P, and A = BC, where k is the number of columns in matrix B. Then, it can be easily verified that $rank_N(A) \le k \le rank(A)$, where $rank_N(A)$ is the non-negative rank of matrix A defined in [3, Definition 7.5.1], and rank(A) is the factor rank of matrix A over the simering A. In this work, we provide some fundamental properties of the A rank of matrices over positive cones of commutative semirings. We also give sufficient conditions for the equality of the A rank, factor rank, and nonnegative rank of matrices. Moreover, we describe the structure of idempotent matrices over positive cones of commutative semirings whose A rank and factor rank coincide.

2. Preliminaries

A semiring [9] is a nonempty set S equipped two binary operations, addition (+) and multiplication (.), such that:

- (S,+) is a commutative monoid with identity element O_S ;
- (S,.) is a monoid with identity element 1_s ;
- Multiplication distributes on both sides with addition;
- $0_S.m = m.0_S = 0_S, \forall m \in S$.

If (S,.) is a commutative monoid then S is a *commutative semiring*. The semiring S is called *zerosumfree* if $a+b=0 \Rightarrow a=b=0, \forall a,b \in S$.

The subset R of S is called a *subsemiring* of the semiring S if $\{0,1\} \subset R$ and it is closed under the addition and multiplication on R.

For two commutative semirings R and S, a mapping $\varphi: S \to R$ is called a *homomorphism* if it satisfies the following conditions:

- i) $\varphi(0_s) = 0_p$, $\varphi(1_s) = 1_p$;
- ii) $\varphi(a+b) = \varphi(a) + \varphi(b), \forall a, b \in S$;
- iii) $\varphi(ab) = \varphi(a)\varphi(b), \forall a, b \in S$.

If there is no risk of confusion, we can write 0 instead of 0_s , 0_R and 1 instead of 1_s , 1_R .

Recall in [4] and [8] that let S be a commutative semiring and P is a subsemiring of S containing the set S^2 , where $S^2 = \{s^2 \mid s \in S\}$. We define $V(P) = \{x \in P \mid \exists y \in P : x + y = 0\}$. If $1 \notin V(P)$ then the subsemiring P is called a *prepositive cone* of the semiring S. Moreover, if the prepositive cone P satisfies the following conditions, it is called a *positive cone* of the semiring S:

- i) If $a \in S$ and $a \notin P$ then there exists an element $b \in P$ such that a+b=0;
- ii) For any $a,b \in S$, if $ab \in V(P)$ then $a \in P$ or $b \in P$.

Notice that, if S is a commutative semiring, then the set P(S) consisting of finite sums of elements in S^2 is a subsemiring of S. Furthermore, if P is a positive cone of S one can have $P(S) \subset P$ (since P is a subsemiring of S and $S^2 \subset P$). In the general case, the subsemiring P(S) may not be a positive cone of the semiring S. According to [8, Theorem 3.3], a commutative semiring S has at least one positive cone if and only if $1+x \neq 0, \forall x \in P(S)$. In that case, we denote PC(S) as the set of positive cones of the semiring S. It is easy to verify that if S is a zero-sumfree semiring, one obtains $PC(S) = \{S\}$.

Example 2.1. Let \mathbb{R} be the semiring of real numbers and \mathbb{B} be the Boolean semiring, we have the set $S = \mathbb{R} \times \mathbb{B}$ forming a commutative semiring with the operations of addition (\oplus) and multiplication (\odot) defined as follows: For any $(a,b),(s,t) \in S$, $(a,b) \oplus (s,t) = (a+s,b+t)$ and $(a,b) \odot (s,t) = (as,bt)$. Here, the additive monoid (S,\oplus) has the unit element (0,0), and the multiplicative monoid (S,\odot) has the unit element (1,1). We have, $V(S) = \{(x,0) \mid x \in \mathbb{R}\}$, this implies that $(1,1) \notin V(S)$. Thus P = S is a positive cone of the semiring S. It is easy to verify that $P(S) = \mathbb{R}^+ \times \mathbb{B}$, where \mathbb{R}^+ is a subsemiring of \mathbb{R} consisting of non-negative real numbers. Since $V(P(S)) = \{(0,0)\}$ hence $(1,1) \notin V(P(S))$, and so, P(S) is a prepositive cone of the semiring S. However, P(S) is not a positive cone of the semiring S since the element (-1,1) does not belong to P(S) and $(-1,1) + (x,y) = (x-1,1) \neq (0,0), \forall (x,y) \in P(S)$.

Example 2.2. For $S = \mathbb{R}^+[x]$, a semiring consisting of polynomials in the variable x with coefficients in \mathbb{R}^+ , we have $P(S) = \left\{ \sum_{i \in J} \left(p_i(x) \right)^2 \mid p_i(x) \in S, 0 < |J| < +\infty \right\}$, where |J| is the number of elements in a certain set J. We have $V(P(S)) = \{0\}$ and $1 \notin V(P(S))$, this follows that P(S) is a prepositive cone of the semiring S. On the other hand, the polynomial $P(x) = x + 1 \in S$ but $P(x) \notin P(S)$, satisfies the condition $P(x) + Q(x) \neq 0, \forall Q(x) \in P(S)$. Therefore, P(S) is not a positive cone of the semiring S.

Proposition 2.3 [8, Lemma 3.5]. Let P be a subset of the commutative semiring S, and $a,b \in S$.

- i) If P is a prepositive cone of S, then $a \in V(P)$ and $a+b \in V(P)$ imply $b \in V(P)$.
- ii) If P is a positive cone of S, then $rm \in V(P)$ for all $m \in V(P)$, $r \in S$.

Note that from Proposition 2.3, we have $ar + bs \in V(P)$ for all $a,b \in V(P)$ and $r,s \in S$. Let S be a commutative semiring, we denote by $M_{m \times n}(S)$ the set of $m \times n$ matrices over semiring S, and $M_n(S)$ is the set of square matrices of order n. The sum of all entries on the main diagonal of matrix $A = (a_{ij}) \in M_n(S)$ is called *trace* of A, denoted by Tr(A). It is easy to verify that for any matrices $A, B \in M_n(S)$, we have Tr(AB) = Tr(BA). A matrix $E \in M_n(S)$ is called idempotent if $E = E^2$. According to [10], two square matrices $E \in M_n(S)$, $F \in M_m(S)$ are said to be equivalent over the semiring S if there exist matrices $A \in M_{n \times m}(S), B \in M_{m \times n}(S)$ such that E = AB and F = BA, and denoted by $E \cong_S F$. Note that, for P being a subsemiring of the semiring S, and $E \in M_n(P), F \in M_m(P)$ being idempotent matrices over P, if $E \cong_P F$ then $E \cong_S F$ but the converse is generally not true. The factor rank of a matrix $A \in M_{m \times n}(S)$ is the smallest non-negative integer k such that there exist matrices $M \in M_{m \times k}(S)$ and $N \in M_{k \times n}(S)$ satisfying A = MN. The factor rank of the matrix A is denoted by rank(A). Note that if $A \in M_{m \times p}(S), B \in M_{p \times n}(S)$ then $rank(AB) \le min\{rank(A), rank(B)\}$. If a matrix A belongs to the set $M_{m \times n}(P(S))$, then A is called a non-negative matrix over the semiring S. According to [3, Definition 7.5.1], non-negative rank of a non-negative matrix $A \in M_{m \times n}(P(S))$, denoted by $rank_N(A)$, is the smallest non-negative integer qsuch that there exist matrices $B \in M_{m \times q}(P(S))$ and $C \in M_{q \times n}(P(S))$ satisfying A = BC. According to [3, Proposition 7.5.2], let S be a commutative semiring, and a matrix $A \in M_{m \times n}(P(S))$, we always have $rank(A) \le rank_N(A) \le min\{m,n\}$. Notice that if $S = \mathbb{R}$, then $P(S) = P(\mathbb{R}) = \mathbb{R}^+$ the semiring of nonnegative real numbers. In [1], Cohen and Rothblum provided a non-negative matrix

$$A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix} \text{ over subsemiring } P(\mathbb{R}) \text{ satisfying the inequality } rank(A) < rank_N(A).$$

Furthermore, by [1, Theorem 4.1], for any non-negative matrix A over subsemiring $P(\mathbb{R})$ of \mathbb{R} , if $rank(A) \leq 2$ then $rank(A) = rank_N(A)$.

3. Main Results

In this section, we investigate some fundamental properties regarding the rank of matrices with entries in a positive cone of a given commutative semiring. Moreover, we compare them with factor rank, non-negative rank, and consider some cases where these rank functions coincide.

Definition 3.1. Let S be a commutative semiring with $PC(S) \neq \emptyset$ and $P \in PC(S)$. The PC rank of a matrix $A \in M_{m \times n}(P)$ is the smallest non-negative integer k such that there exist matrices $B \in M_{m \times k}(P)$ and $C \in M_{k \times n}(P)$ satisfying A = BC. The PC rank of matrix $A \in M_{m \times n}(P)$ is denoted by $rank_{PC}(A)$.

Remark 3.2. If P is a positive cone of the commutative semiring S, then for any $A \in M_{m \times n}(P)$, we always have $A = AI_n = I_m A$. Since entries of matrices I_m, I_n belong to P, the PC rank of matrix A always exists and satisfies the condition $rank_{PC}(A) \le \min\{m,n\}$. In addition, since $P(S) \subset P \subset S$, hence, for any matrix $A \in M_{m \times n}(P(S))$, we get the inequality $rank(A) \le rank_{PC}(A) \le rank_N(A)$. The following examples provide cases where the equal sign "=" does not hold.

Example 3.3. Consider the commutative semiring $S = \mathbb{R}$ with $P(S) = \mathbb{R}^+$ and $V(P(S)) = \{0\}$. Therefore, $1 \notin V(P(S))$. For any $x \in S$ and $x \notin P(S)$, there always exists an element $-x \in P(S)$ such that x + (-x) = 0. Furthermore, for any $x, y \in S$ such that $x, y \in V(P(S)) = \{0\}$, we have $x = 0 \in P(S)$ or $y = 0 \in P(S)$. Thus P = P(S) is a positive cone of S. We have the matrix

$$A = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} \in M_4(P(S)). \text{ According to [1], } rank(A) = 3 < 4 = rank_N(A). \text{ Moreover, since}$$

P = P(S) hence $rank_N(A) = rank_{PC}(A)$, and so $rank(A) < rank_{PC}(A) = rank_N(A)$.

Lemma 3.4. Let S,R be commutative semirings. If there exists a homomorphism $\varphi:S\to R$, then for any matrix $A=(a_{ij})\in M_{m\times n}\left(P(S)\right)$, we have the matrix $\varphi(A)$ defined by $\varphi(A)=\left(\varphi(a_{ij})\right)\in M_{m\times n}\left(P(R)\right)$, satisfying the inequality $rank_N\left(\varphi(A)\right)\leq rank_N\left(A\right)$.

Proof. For any matrix $A = (a_{ij}) \in M_{m \times n}(P(S))$, we have $a_{ij} \in P(S), \forall i = 1, ..., m; j = 1, ..., n$. Since φ is a homomorphism hence $\varphi(a_{ij}) \in P(R), \forall i = 1, ..., m; j = 1, ..., n$, this follows that $\varphi(A) = (\varphi(a_{ij})) \in M_{m \times n}(P(R))$. Suppose that $rank_N(A) = k$, then there exist matrices $B \in M_{m \times k}(P(S)), C \in M_{k \times n}(P(S))$ such that A = BC, and so $\varphi(A) = \varphi(B).\varphi(C)$ (because φ is a homomorphism). This implies that $rank_N(\varphi(A)) \le k = rank_N(A)$. \square

Example 3.5. Consider the commutative semiring $S = \mathbb{R} \times \mathbb{B}$ introduced in section 2 and the

matrix
$$A = \begin{pmatrix} (1,0) & (0,0) & (0,0) & (1,0) \\ (1,0) & (1,0) & (0,0) & (0,0) \\ (0,0) & (1,0) & (1,0) & (0,0) \\ (0,0) & (0,0) & (1,0) & (1,0) \end{pmatrix}$$
, we observe that entries of A belong to $P(S) = \mathbb{R}^+ \times \mathbb{B}$.

Then, P = S is a positive cone of the semiring S. Furthermore, we have:

$$A = \begin{pmatrix} (1,0) & (0,0) & (0,0) \\ (1,0) & (1,0) & (0,0) \\ (0,0) & (1,0) & (1,0) \\ (0,0) & (0,0) & (1,0) \end{pmatrix} \begin{pmatrix} (1,1) & (0,0) & (0,0) & (1,0) \\ (0,0) & (1,1) & (0,0) & (-1,0) \\ (0,0) & (0,0) & (1,1) & (1,0) \end{pmatrix}, \text{ this follows}$$
 that:
$$rank(A) = rank_{PC}(A) \le 3.$$

Now, consider the mapping $\varphi: S \to \mathbb{R}$ defined by $\varphi((x,y)) = x, \forall (x,y) \in S$. It is easy to verify

that φ is a homomorphism and $\varphi(A) = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} \in M_4(P(\mathbb{R})) = M_4(\mathbb{R}^+)$. Applying Lemma 3.4,

we obtain the inequality $rank_N(\varphi(A)) \le rank_N(A)$. By Example 3.3, we have $rank_N(\varphi(A)) = 4$, and so $rank_N(A) = 4$. Thus $rank_{PC}(A) \le 3 < 4 = rank_N(A)$.

Next, we provide some basic properties regarding the PC rank of matrices over the positive cones of a given commutative semiring. The following proposition provides some basic properties of the PC rank, and the proof is similar to the one introduced in [11].

Proposition 3.6. Let P is a positive cone of the commutative semiring S and matrices $A \in M_{m \times n}(P)$, $B \in M_{n \times p}(P)$, $C \in M_{m \times q}(P)$, $D \in M_{m \times n}(P)$, $E \in M_{k \times l}(P)$. Then, the following statements hold:

- i) $rank_{PC}(AB) \le min\{rank_{PC}(A), rank_{PC}(B)\}.$
- ii) $\max \{ rank_{PC}(A), rank_{PC}(C) \} \le rank_{PC}(A C) \le rank_{PC}(A) + rank_{PC}(C)$.

$$iii) \ \max \left\{ rank_{PC} \left(A \right), rank_{PC} \left(E \right) \right\} \leq rank_{PC} \left(\begin{matrix} A & 0 \\ 0 & E \end{matrix} \right) \leq rank_{PC} \left(A \right) + rank_{PC} \left(E \right).$$

iv) $rank_{PC}(A+D) \le min\{rank_{PC}(A) + rank_{PC}(D), m, n\}.$

Where $\begin{pmatrix} A & C \end{pmatrix}$ and $\begin{pmatrix} A & 0 \\ 0 & E \end{pmatrix}$ are block matrices formed by matrices A, C, E.

Proof.

Suppose that $rank_{PC}\left(A\right) = r, rank_{PC}\left(B\right) = s, rank_{PC}\left(C\right) = u$, then there exist matrices $M \in M_{m \times r}\left(P\right), \ N \in M_{r \times n}\left(P\right), \ H \in M_{n \times s}\left(P\right), \ G \in M_{s \times p}\left(P\right), \ X \in M_{m \times u}\left(P\right) \ \text{and} \ Y \in M_{u \times q}\left(P\right) \ \text{such that} \ A = MN \ , \ B = HG \ \text{and} \ C = XY \ .$

 $i) \text{ Since } AB = MNHG \text{ , hence, } rank_{PC} \left(AB \right) \leq \min \left\{ r, s \right\} = \min \left\{ rank_{PC} \left(A \right), rank_{PC} \left(B \right) \right\}.$

ii) We have
$$(A \ C) = (MN \ XY) = (M \ X) \begin{pmatrix} N & 0 \\ 0 & Y \end{pmatrix}$$
, this implies that $rank_{PC}(A \ C) \le r + u = rank_{PC}(A) + rank_{PC}(C)$.

Suppose that $rank_{PC}(A \ C) = v$, then there exist matrices $M' \in M_{m \times v}(P), N' \in M_{v \times (n+q)}(P)$ such that $(A \ C) = M'N'$. Set $N' = (H' \ G')$, where $H' \in M_{v \times n}(P), G' \in M_{v \times q}(P)$. We have $(A \ C) = (M'H' \ M'G')$, hence, A = M'H', C = M'G'. Thus $rank_{PC}(A) \le v$ and $rank_{PC}(C) \le v$, this implies that $\max \{rank_{PC}(A), rank_{PC}(C)\} \le v = rank_{PC}(A \ C)$.

iii) and iv) follow from i) and ii).

Definition 3.7. Let P be a positive cone of the commutative semiring S. A square matrix $E \in M_n(P)$ is called PC - full if $rank_{PC}(E) = n$.

Remark 3.8. If P is a positive cone of the commutative semiring S, then every identity matrix I_n is PC - full. Indeed, since S is a commutative semiring, hence, P is also a commutative semiring. For any matrices $A, B \in M_k(P)$ such that $AB = I_k$, then by [12], we have $BA = I_k$. Applying [10, Proposition 3.12] to the semiring P with PC rank, we obtain the following result: For any matrices $G \in M_{n \times m}(P), H \in M_{m \times n}(P)$ such that $GH = I_n$, it follows that $m \ge n$. Thus $rank_{PC}(I_n) = n$. Similarly, applying [10, Lemma 3.12] to the commutative semiring S, we also have $rank(I_n) = n$.

Note that, for any matrix $A \in M_{m \times n}(P)$, where P is a positive cone of the commutative semiring S, if $rank_{PC}(A) = k$ then there exist matrices $B \in M_{m \times k}(P)$ and $C \in M_{k \times n}(P)$ such that $A = BC = BI_kC$. This gives us a hint on how to compute the PC rank of matrices through the PC rank of PC - full idempotent matrices, as showed in the following proposition.

Proposition 3.9. Let P be a positive cone of the commutative semiring S and a matrix $A \in M_{m \times n}(P)$. Then, $rank_{PC}(A) = \min\{rank_{PC}(E) | A = BEC, E \in \mathbb{EM}_{fpc}(P)\}$, where B, C are matrices with entries in P, $\mathbb{EM}_{fpc}(P)$ is the set of PCs - full idempotent matrices over P.

Proof. We have $A = I_m I_m A = A I_n I_n$, this implies that the set $\{rank_{PC}(E) | A = BEC, E \in \mathbb{EM}_{fpc}(P)\}$ is a nonempty set, where B, C are matrices with entries in P. Furthermore, if $rank_{PC}(A) = k$ then there exist matrices $B \in M_{m \times k}(P)$ and $C \in M_{k \times n}(P)$ such that $A = BC = BI_k C$, since $rank_{PC}(I_k) = k$ (by Remark 3.8), hence, we obtain the inequality $rank_{PC}(A) \ge \min \{rank_{PC}(E) | A = BEC, E \in \mathbb{EM}_{fpc}(P)\}$. Conversely, suppose that $\min \{rank_{PC}(E) | A = BEC, E \in \mathbb{EM}_{fpc}(P)\} = k$, then there exists a matrix $E \in \mathbb{EM}_{fpc}(P)$ with order k, and matrices $B \in M_{m \times k}(P), C \in M_{k \times n}(P)$ such that A = BEC and $rank_{PC}(E) = k$. Applying Proposition 3.6, we have $rank_{PC}(A) = rank_{PC}(BEC) \le rank_{PC}(E) = k$. So, the proposition has been proven. □

Next, we investigate some conditions under which the PC rank of matrices coincides with the non-negative rank or the factor rank.

Proposition 3.10. Let P be a positive cone of the commutative semiring S. Then,

- i) If $A \in M_n(P)$ is an invertible matrix over the semiring S then $rank_{PC}(A) = rank(A) = n$.
- ii) If $A \in M_n(P(S))$ is an invertible matrix over the semiring S then $rank_N(A) = rank_{PC}(A) = rank(A) = n$.

Proof.

i) If $A \in M_n(P)$ with $rank(A) = k \le n$, then there exist matrices $B \in M_{n \times k}(S)$ và $C \in M_{k \times n}(S)$ such that A = BC. Because A is an invertible matrix over the semiring S, hence, $I_n = A^{-1}BC$. By Remark 3.8, we have $n = rank(I_n) \le k$, and so rank(A) = n. Moreover, $rank(A) \le rank_{PC}(A) \le n$. Thus $rank_{PC}(A) = rank(A) = n$.

ii) If an invertible matrix $A \in M_n(P(S))$ with $rank(A) = k \le n$, similarly, we can show that rank(A) = n. Furthermore, $n = rank(A) \le rank_{PC}(A) \le rank_N(A) \le n$, this follows that $rank_N(A) = rank_{PC}(A) = rank(A) = n$. \square

Remark 3.11. Let P be a positive cone of the commutative semiring S, if $A \in M_n(P(S))$ is an invertible matrix over the semiring S, then A^{-1} may not belong to the set $M_n(P(S))$. Moreover, if $A \in M_n(P)$ is an invertible matrix over the semiring S, then A^{-1} may not belong to the set $M_n(P)$. For example, consider the semiring $S = \mathbb{R}$ consisting of real numbers, we have $A = \begin{pmatrix} 1 & 1 \\ 0 & 4 \end{pmatrix} \in M_2(P(\mathbb{R}))$ is an invertible matrix over \mathbb{R} , and $A^{-1} = \begin{pmatrix} 1 & -\frac{1}{4} \\ 0 & \frac{1}{4} \end{pmatrix} \notin M_2(P(\mathbb{R}))$. It is easy to

verify that $P(\mathbb{R}) = \mathbb{R}^+$, and $P = P(\mathbb{R})$ is a positive cone of S. Then, $A^{-1} \notin M_2(P)$. However,

consider the matrix
$$B = \begin{pmatrix} 0 & 4 \\ 9 & 0 \end{pmatrix} \in M_2(P(\mathbb{R}))$$
, we have $B^{-1} = \begin{pmatrix} 0 & \frac{1}{9} \\ \frac{1}{4} & 0 \end{pmatrix} \in M_2(P(\mathbb{R}))$.

Proposition 3.12. Let P be a positive cone of the commutative semiring S, and Q be a subsemiring of S such that $P \subset Q$. Then, the following statements are equivalent:

- i) If any square matrix $A \in M(P)$ is invertible over Q, then $A^{-1} \in M(P)$.
- ii) P = Q.

Proof.

- $(ii) \Rightarrow i$): Obviously.
- $i)\Rightarrow ii)$: If $P\subset Q$ and $P\neq Q$, then there exists an element $a\in Q\setminus P$. Since P is a positive cone of the semiring S, hence, there exists an element $b\in P$ such that a+b=0. We have $A=\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\in M_2(P)$ and $\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}=\begin{pmatrix} 1 & a+b \\ 0 & 1 \end{pmatrix}=\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, this implies that $A^{-1}=\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}\in M_2(Q)$. So, the matrix A is invertible over Q. Therefore, $A^{-1}\in M_2(P)$, and so $a\in P$, this is contradictory to $a\notin P$. Thus P=Q. \square

The following Corollary can be readily deduced from the Proposition 3.12.

Corollary 3.13. Let P is a positive cone of the commutative semiring S. Then, the following statements hold:

i) If P(S) is a positive cone of the semiring S, and if any square matrix A in M(P(S)), which is invertible over the semiring P, always has its inverse A^{-1} also in M(P(S)), then, $rank_N(B) = rank_{PC}(B), \forall B \in M(P(S))$.

- *ii*) If any square matrix A in M(P), which is invertible over the semiring S, always has its inverse A^{-1} also in M(P), then, $rank(B) = rank_{PC}(B)$, $\forall B \in M(P)$.
- *iii)* If P(S) is a positive cone of the semiring S, and if any square matrix A in M(P(S)), which is invertible over the semiring S, always has its inverse A^{-1} also in M(P(S)), then, $rank_N(B) = rank_{PC}(B) = rank(B), \forall B \in M(P(S))$.

Next, we examine the cases where the PC rank and the factor rank of idempotent matrices are equal over commutative semirings.

Theorem 3.14. Let P is a positive cone of the commutative semiring S, and $E \in M_n(V(P))$ be an idempotent matrix. Then, $rank(E) = rank_{PC}(E)$.

Proof. Suppose that rank(E) = k, then there exist matrices $B \in M_{n \times k}(S)$ and $C \in M_{k \times n}(S)$ such that E = BC. Let $F = CEB \in M_k(S)$, we have $F^2 = (CEB)(CEB) = CE^3B = CEB = F$, this follows that F is an idempotent matrix. Furthermore, since $E \in M_n(V(P))$, hence, by Proposition 2.3, we have $CE \in M_{k \times n}(V(P))$, $EB \in M_{n \times k}(V(P))$, and so $F = (CE)(EB) \in M_k(V(P))$, this implies that $BF \in M_{n \times k}(V(P))$, $FC \in M_{k \times n}(V(P))$. Moreover, we can decompose the matrix E into $E = E^5 = B(CBCB)(CBCB)C = (BF)(FC)$, hence, $rank_{PC}(E) \le k = rank(E)$. Therefore, $rank(E) = rank_{PC}(E)$. \Box

Corollary 3.15. Let P is a positive cone of the commutative semiring S, and $E \in M_n(P)$ be an idempotent matrix with $Tr(E) \in V(P)$. Then, the following statements hold:

- i) If rank(E)=1 then $rank_{PC}(E)=1$.
- ii) If rank(E) = 2, then there exists an idempotent matrix $F \in M_2(P)$ that is PC full and $E \cong_S F$, such that $rank(E) = rank(F) = rank_{PC}(F) = 2$.

Proof.

- i) If rank(E) = 1, then there exist matrices $B = (b_{i1}) \in M_{n \times 1}(S)$ and $C = (c_{1j}) \in M_{1 \times n}(S)$ such that E = BC. Then, $Tr(E) = \sum_{l=1}^{n} c_{1l}b_{l1} \in V(P)$, this follows that $F = CB \in M_1(V(P))$. Hence, $BF \in M_{n \times 1}(V(P))$ and $FC \in M_{1 \times n}(V(P))$, and so $E = E^3 = (BF)(FC) \in M_n(V(P))$. By Theorem 3.14, we obtain $rank_{PC}(E) = rank(E) = 1$.
- ii) If rank(E) = 2, then there exist matrices $B = (b_{ij}) \in M_{n \times 2}(S)$ and $C = (c_{ij}) \in M_{2 \times n}(S)$ such that E = BC. Let $H = CB = \begin{pmatrix} q_1 & p_1 \\ p_2 & q_2 \end{pmatrix} \in M_2(S)$, we have $q_1 + q_2 = Tr(CB) = Tr(BC) = Tr(E) \in V(P)$

and
$$H^2 = \begin{pmatrix} q_1^2 + p_1 p_2 & p_1(q_1 + q_2) \\ p_2(q_1 + q_2) & q_2^2 + p_1 p_2 \end{pmatrix} = CEB$$
. Setting $F = H^2$, we have

 $F^2 = (CEB)(CEB) = CE^3B = CEB = F$, hence, F is an idempotent matrix and

$$F = F^{2} = \begin{pmatrix} \left(q_{1}^{2} + p_{1}p_{2}\right)^{2} + p_{1}p_{2}\left(q_{1} + q_{2}\right)^{2} & \left(q_{1} + q_{2}\right)\left[p_{1}\left(q_{1}^{2} + p_{1}p_{2}\right) + p_{1}\left(q_{2}^{2} + p_{1}p_{2}\right)\right] \\ \left(q_{1} + q_{2}\right)\left[p_{2}\left(q_{1}^{2} + p_{1}p_{2}\right) + p_{2}\left(q_{2}^{2} + p_{1}p_{2}\right)\right] & \left(q_{2}^{2} + p_{1}p_{2}\right)^{2} + p_{1}p_{2}\left(q_{1} + q_{2}\right)^{2} \end{pmatrix}.$$

Because $\left(q_1^2+p_1p_2\right)^2$, $\left(q_2^2+p_1p_2\right)^2\in P(S)\subset P$ and $q_1+q_2\in V(P)$, applying Proposition 2.3, we have the elements $p_1p_2\left(q_1+q_2\right)^2$, $\left(q_1+q_2\right)\left[p_1\left(q_1^2+p_1p_2\right)+p_1\left(q_2^2+p_1p_2\right)\right]$ and $\left(q_1+q_2\right)\left[p_2\left(q_1^2+p_1p_2\right)+p_2\left(q_2^2+p_1p_2\right)\right]$ belonging to $V(P)\subset P$. Thus the idempotent matrix $F\in M_2\left(P\right)$. Moreover, since $E=E^3=BFC$, hence, $2=rank(E)\leq rank(F)\leq rank_{PC}(F)\leq 2$, this follows that $rank_{PC}\left(F\right)=rank\left(F\right)=rank\left(E\right)=2$, and so F is a PC - full matrix. In additon, because $(EB)C=E^2=E$ and C(EB)=F, hence, $E\cong_S F$. \square

Remark 3.16. If the matrix $F = H^2 = \begin{pmatrix} q_1^2 + p_1 p_2 & p_1 (q_1 + q_2) \\ p_2 (q_1 + q_2) & q_2^2 + p_1 p_2 \end{pmatrix}$, as determined in Corollary 3.15, has all its entries on the main diagonal belonging to V(P), then $F \in M_2(V(P))$. This follows that $E = BFC = (BF)(FC) \in M_n(V(P))$. By Theorem 3.14, we obtain the equality $rank(E) = rank_{PC}(E) = rank(F) = rank_{PC}(F) = 2$.

Example 3.17. The semiring $S = \mathbb{R} \times \mathbb{B}$ has $V(S) = \{(x,0) | x \in \mathbb{R}\}$, hence, $(1,1) \notin V(S)$. We have

$$P = S \text{ as a positive cone of the semiring } S. \text{ Then, } E = \begin{pmatrix} \left(\frac{1}{2}, 0\right) & (-2, 0) \\ \left(-\frac{1}{8}, 0\right) & \left(\frac{1}{2}, 0\right) \end{pmatrix}, G = \begin{pmatrix} \left(\frac{1}{2}, 0\right) & (1, 0) & (0, 0) \\ \left(\frac{1}{4}, 0\right) & \left(\frac{1}{2}, 0\right) & (0, 0) \\ (0, 0) & (0, 0) & (1, 0) \end{pmatrix}$$

are idempotent matrices over the positive cone P. Furthermore, $Tr(E) = (1,0) \in V(P)$, $Tr(G) = (2,0) \in V(P)$ and rank(E) = 1, rank(G) = 2.

Lemma 3.18. Let S be a commutative semiring, and $E = (e_{ij}) \in M_n(S)$ be an square matrix. Then for any $k \in \{1, 2, ..., n\}$, the entries of the matrix $B = E^k$ are determined by the formula $b_{ij} = \begin{cases} \sum_{l_1, l_2, ..., l_{k-1} \in \{1, 2, ..., n\}} e_{il_1} e_{l_1 l_2} ... e_{l_{k-2} l_{k-1}} e_{l_{k-1} j}, \ k > 1 \\ e_{ii}, k = 1 \end{cases}, \forall i, j \in \{1, 2, ..., n\}, \text{ where } B = (b_{ij}).$

 $Proof. \ \, \text{If} \ \ \, k=1 \ \, \text{then} \ \, B=E \, , \, \, \text{hence}, \quad b_{ij}=e_{ij}, \forall i,j\in\left\{1,2,...,n\right\}. \, \, \text{If} \ \ \, k=2 \ \, \text{then} \ \, B=E^2 \, , \, \, \text{hence}, \\ b_{ij}=\sum_{l\in\left\{1,2,...,n\right\}}e_{il}e_{lj}, \forall i,j\in\left\{1,2,...,n\right\}. \, \, \text{Suppose that the Lemma is true with} \, \, k=q, \, \left(2\leq q < n\right). \, \, \text{Let} \, \, C=E^q \, , \\ \text{the entries of the matrix} \, \, B=E^{q+1}=CE \, \, \text{are determined by} \, \, b_{ij}=\sum_{t\in\left\{1,2,...,n\right\}}c_{it}e_{ij} \, , \, \, \text{where} \, \, C=\left(c_{ij}\right) \, \, \text{and} \\ B=\left(b_{ij}\right). \, \, \text{Since} \, \, c_{ii}=\sum_{l_1,l_2,...,l_{q-1}\in\left\{1,2,...,n\right\}}e_{il_1}e_{l_1l_2}...e_{l_{q-2}l_{q-1}}\, e_{l_{q-1}t} \, e_{l_{q-1}t} \, , \, \, \text{hence}, \, \, b_{ij}=\sum_{l_1,l_2,...,l_{q-1},t\in\left\{1,2,...,n\right\}}e_{il_1}e_{l_1l_2}...e_{l_{q-2}l_{q-1}}\, e_{l_{q-1}t} \, e_{ij} \, . \\ \text{the Lemma has been proven.} \, \, \Box$

Lemma 3.19. Let *P* be a positive cone of the commutative semiring *S*, and $A = (a_{ij}) \in M_n(P)$ be an idempotent matrix. If $a_{ii} = 0, \forall i \in \{1,...,n\}$ then $A = M_n(V(P))$.

 $Proof. \ \ If \ n=1 \ \ \text{then} \ \ A=\left(0\right)\in M_1\left(V\left(P\right)\right). \ \ If \ n>1 \ , \ \text{then for each} \ \ k\in\left\{1,2,...,n\right\}, \ \ \text{by Lemma 3.18,}$ the entries of the main diagonal of the matrix $B=A^k$ are determined by the formula $b_{ii}=\begin{cases} \sum_{l_1,l_2,...,l_{k-1}\in\{1,2,...,n\}} a_{il_1}a_{l_1l_2}...a_{l_{k-1}i}, \ k>1\\ 0,\ k=1 \end{cases}, \ \forall i\in\left\{1,2,...,n\right\}, \ \ \text{where} \ \ B=\left(b_{ij}\right). \ \ \text{Since} \ \ A=\left(a_{ij}\right) \ \ \text{is an} \end{cases}$

 $a_{i l_1} a_{l_1 l_2} \dots a_{l_{k-2} l_{k-1}} a_{l_{k-1} i} \in V\left(P\right), \forall i \in \left\{1, 2, \dots, n\right\} \text{ (because } A = \left(a_{i j}\right) \in M_n\left(P\right) \text{). Now let } E = A^n \text{, then the entries of the matrix } E \text{ are determined by the following formula: } \\ e_{i j} = \begin{cases} \sum\limits_{l_1, l_2, \dots, l_{n-1} \in \left\{1, 2, \dots, n\right\}} a_{i l_1} a_{l_1 l_2} \dots a_{l_{n-2} l_{n-1}} a_{l_{n-1} j}, \ i \neq j \\ 0, \ i = j \end{cases}, \forall i, j \in \left\{1, 2, \dots, n\right\}. \text{ Let } l_0 = i, \ l_n = j \text{, then for any late } l_n = j \text{, then late } l_n = j \text{,$

$$\begin{split} &l_{1},l_{2},...,l_{n-1}\in\left\{ 1,2,...,n\right\} ,\text{ since }l_{0},l_{1},...,l_{n}\in\left\{ 1,2,...,n\right\} ,\text{ hence, there exist at least two elements }\\ &r,t\in\left\{ 0,1,...,n\right\} \text{ such that }r< t\text{ and }l_{r}=l_{t}.\text{ This follows that }a_{l_{r}l_{r+1}}...a_{l_{t-1}l_{t}}\in V\left(P\right),\text{ and so }\\ &a_{il_{1}}a_{l_{i}l_{2}}...a_{l_{n-2}l_{n-1}}a_{l_{n-1}j}=a_{il_{1}}a_{l_{i}l_{2}}...a_{l_{r-1}l_{t}}...a_{l_{r-2}l_{n-1}}a_{l_{n-1}j}\in V\left(P\right) \text{ (by Proposition 2.3), therefore, }\\ &e_{ii}\in V\left(P\right),\forall i,j\in\left\{ 1,2,...,n\right\} .\text{ Thus }E=A^{n}=A\in M_{n}\left(V\left(P\right)\right).\ \Box \end{split}$$

Corollary 3.20. Let S be a commutative ring, and Q be a positive cone of the commutative semiring R. Then, the following statements hold:

- i) The semiring $P = S \times Q$ is a positive cone of the semiring $S \times R$.
- ii) If $E \in M_n(P)$ is an idempotent matrix with entries on the main diagonal belonging to set $S \times \{0_R\}$, then $rank(E) = rank_{PC}(E)$.

Proof.

i) Since Q is a positive cone of the commutative semiring R, hence, $P = S \times Q$ is a subsemiring of the semiring $S \times R$. It easy to see that $V(P) = S \times V(Q)$ (because S is a commutative ring). We have $(1_S, 1_R) \notin V(P)$ (since Q is a positive cone of R hence $1_R \notin V(Q)$). This follows that P is a prepositive

cone of the semiring $S \times R$. For any element $(x,y) \in S \times R$ and $(x,y) \notin P = S \times Q$, we have $y \notin Q$. Since Q is a positive cone of R, hence, there exists an element $b \in Q$ such that $y+b=0_R$, therefore, the element $(-x,b) \in P$ satisfies $(x,y)+(-x,b)=(0_S,0_R)$. Furthermore, if there exist elements $(x,y),(u,v) \in S \times R$ such that $(x,y)(u,v)=(xu,yv) \in S \times V(Q)$, then $yv \in V(Q)$, and so, $y \in Q$ or $v \in Q$. This implies that $(x,y) \in S \times Q = P$ or $(u,v) \in S \times Q = P$. Thus P is a positive cone of the semiring $S \times R$.

ii) If the idempotent matrix $E = (e_{ij}) \in M_n(P)$ has entries on the main diagonal belonging to the set $S \times \{0_R\}$, then let $e_{ij} = (s_{ij}, q_{ij}), \forall i, j \in \{1, ..., n\}$, we have $e_{ii} = (s_{ii}, 0_R), \forall i \in \{1, ..., n\}$. Consider the homomorphism $\varphi \colon S \times R \to R$ with $\varphi(s,q) = q, \forall (s,q) \in S \times R$. Then, the matrix $\varphi(E) = (q_{ij}) \in M_n(Q)$ is an idempotent matrix with entries on the main diagonal equal to 0_R . By Lemma 3.19, we have $\varphi(E) \in M_n(V(Q))$, this follows that $e_{ij} = (s_{ij}, q_{ij}) \in S \times V(Q) = V(P), \forall i, j \in \{1, ..., n\}$, and so $E \in M_n(V(P))$. Applying Theorem 3.14, we obtain $rank(E) = rank_{PC}(E)$. \square

Acknowledgments

The authors express their deep gratitude to the anonymous referee for the extremely careful reading and highly professional working with our manuscript.

References

- [1] J. E. Cohen, U. G. Rothblum, Nonnegative Ranks, Decompositions, and Factorizations of Nonnegative Matrices, Linear Algebra Appl., Vol. 190, 1993, pp. 149-168.
- [2] L. B. Beasley, T. J. Laffey, Real Rank Versus Nonnegative Rank, Linear Algebra Appl., Vol. 431, No. 12, 2009, pp. 2330-2335, https://doi.org/10.1016/j.laa.2009.02.034.
- [3] P. Mohindru, Completely Positive Matrices Over Semirings and Their CP-rank, Diss. University of Guelph, 2014.
- [4] P. Mohindru, Pereira, Orderings on Semirings and Completely Positive Matrices, Linear Multilinear Algebr., Vol. 64, No. 5, 2016, pp. 818-833, https://doi.org/10.1080/03081087.2015.1059405.
- [5] Y. Shitov, On Tropical and Nonnegative Factorization Ranks of Band Matrices, arXiv:1710.02072v1, 2017.
- [6] D. K. Bukovšek, H. Šmigoc, Symmetric Nonnegative Matrix Trifactorization, Linear Algebra Appl., Vol. 665, 2023, pp. 36-60, https://doi.org/10.1016/j.laa.2023.01.027.
- [7] L. Q. Tung, T. Hoang, Asymptotic Nonnegative Rank of Matrices, arXiv:2308.07187, 2023.
- [8] T. C. Craven, Orderings on Semirings, Semigr, Forum, Vol. 43, No. 1, 1991, pp. 45-52, https//doi.org/10.1007/BF02574250.
- [9] J. S. Golan, Semirings and their Applications. Kluwer Academic Publishers, Dordrecht-Boston-London, 1999.
- [10] H. C. Cong, Stably Free Rank of Idempotent Matrices on Semirings, Univ. Da Nang, J. Sci. Technol., Vol. 20, 2022, pp. 56-60.
- [11] L. B. Beasley, A. E. Guterman, Rank Inequalities Over Semirings, J. Korean Math. Soc., Vol. 42, No. 2, 2005, pp. 223-241.
- [12] C. Reutenauer, H. Straubing, Inversion of Matrices Over A Commutative Semiring, J. Algebr, Vol. 88, No. 2, 1984, pp. 350-360.