

# Chemical composition of essential oil of *Machilus Japonica* Siebold & Zucc. (Lauraceae) from Vietnam

Pham Van Khien<sup>1</sup>, Do Quang Huy<sup>2,\*</sup>, Do Thi Viet Huong<sup>2</sup>

<sup>1</sup>*Pharmaceutical Journal, Vietnam*

<sup>2</sup>*College of Science, VNU, 334 Nguyen Trai, Hanoi, Vietnam*

Received 29 October 2008

**Abstract.** Chemical composition of essential oil of *Machilus Japonica* Siebold & Zucc. (Lauraceae) from Vietnam were studied. The essential oil were obtained from the leaves of young plants (3-4 years) by hydrodistillation. The identification of the components was performed using GC, GC/MS. The main components in this species identified were  $\alpha$ -phellandrene (60.2%),  $\beta$ -phellandrene (10.5%), p-cymene (20.3%),  $\beta$ -myrcene (3.7%), terpineole (0.9%), trans- $\beta$ -ocimene (0.7%), 11-dodecenol (0.3),  $\beta$ -Caryophyllene (0,5%). This result were not reported in any of previously investigated species of Lauraceae family in Vietnam. This was a new species of Lauraceae family in Vietnam. The name of this species was *Machilus japonica* Siebold & Zucc.

**Keywords:** Lauraceae; *Machilus japonica*; leaves; essential oil; phellandrene; p-cymene.

## Introduction

Lauraceae is lager family that includes 32 genera and more than 2500 species. it is known about 256 species (21 genera) in Vietnam has been described. In this paper, the chemical composition of essential oil of *Machilus Japonica* Siebold & Zucc. (Lauraceae) from Vietnam will be discussed. The essential oil composition of the different species of Lauraceae family in Vietnam have been reported by many authors [1-8]. These varieties are morphologically similar, but they show different chemical compositions of essential oil. Therefore, the chemical composition of

essential oil can be use to identify the species of Lauraceae family.

## Materials and methods

**Plant material.** The leaves of young plants of Lauraceae family (3-4 years) were collected in October 1991 and 2007 at around Ba Vi mountain, in Northern Vietnam. The essential oils of leaves were obtained with a yield of 1.23% (w/w) by a 1 hour hydrodistillation in a Clevenger-type apparatus, and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>.

**Analytical.** GC analysis were performed on a Hewlett Packard 5890 Series equipped with a split/splitless injector; flame ionization detector (FID); capillary column HP-5 (30 m x 320  $\mu$ m

\* Corresponding author. Tel.: 84-4-8583905  
E-mail: [huydq@vnu.edu.vn](mailto:huydq@vnu.edu.vn)

i.d., film thickness 0.25  $\mu\text{m}$ ); column temperature, 60°C for 3 min, 5°C/min to 230°C and kept at 230°C for 10 min; injector temperature 250°C; split ratio 1:40; detector temperature 260°C; carrier gas N<sub>2</sub> 0.9 ml/min; volume injected 1  $\mu\text{L}$ . GC/MS analysis was performed on a Hewlett Packard 5890 Series equipped with HP5971A mass selective detector; capillary column HP-5MS (30 m x 250  $\mu\text{m}$  i.d., film thickness 0.25  $\mu\text{m}$ ); column temperature, 60°C for 3 min, 5°C/min to 230°C and kept at 230°C for 10 min; injector temperature 250°C; split ratio 1:50; detector temperature 260°C; carrier gas He 0.9 ml/min; volume injected 1  $\mu\text{L}$ ; interface temperature 280°C; acquisition mass range 40-400. Identification of the constituents of essential oil was made by comparison of their mass spectra with nbs751 library spectra data. Relative percentage amounts were calculated from FID peak areas [9-12].

## Results and discussion

The essential oil was light yellow and the total yield of 1.23%. The chemical composition of the leaf essential oils of investigated species collected in October 1991 and 2007 were not different and included 18 components (table 1). As it is shown in table 1, the leaf essential oil of investigated species from Vietnam was characterized by large amount of  $\alpha$ - and  $\beta$ -phellandrene (70.7%) and the ratio of  $\alpha$ -phellandrene and  $\beta$ -phellandrene was 5.73.

Table 1. The composition of leaf essential oil of *Machilus japonica* Siebold & Zucc. (Lauraceae) from Vietnam

Peak	Identification	KI	Percent
1	$\alpha$ -Thujene	931	trace
2	$\alpha$ -Pinene	939	0.2
3	Sabinene	976	trace

4	$\beta$ -Pinene	980	trace
5	$\beta$ -Myrcene	990	3.7
6	$\alpha$ -Phellandrene	1006	60.2
7	p-Cymene	1028	20.3
8	$\beta$ -Phellandrene	1031	10.5
9	cis- $\beta$ -Ocimene	1042	0.2
10	trans- $\beta$ -Ocimene	1053	0.7
11	Linalool	1100	trace
12	Camphor	1145	trace
13	$\alpha$ -Terpineol	1189	0.9
14	$\alpha$ -Phellandrene ocide	1192	0.1
15	(Z)- $\beta$ -Carvotancetol	1210	0.1
16	11-Dodecenol	1369	0.3
17	Carvarol	1371	0.2
18	$\beta$ -Caryophyllene	1414	0.5
Total			97.8

The main components in the leaf essential oil were  $\alpha$ -phellandrene (60.2%),  $\beta$ -phellandrene (10.5%), p-cymene (20.3%). Other notable constituents were  $\beta$ -myrcene (3.7%),  $\alpha$ -terpineol (0.9%), trans- $\beta$ -ocimene (0.7%), 11-Dodecenol (0.3),  $\beta$ -Caryophyllene (0.5%). The high content of  $\alpha$ -phellandrene,  $\beta$ -phellandrene and p-cymene were detected in the leaf essential oil, which were not reported in any of previously investigated species of Lauraceae family in Vietnam. To the best of our knowledge, this was a new species of Lauraceae family in Vietnam. The name of this species was *Machilus japonica* Siebold & Zucc.

## Acknowledgements

The authors are grateful for financial support by National project for base research from the Ministry of Science and Technology of Vietnam, code 50 63 06.

## References

- [1] N.X. Dung, P.V. Khien, Essential oils of wood, root, flower, and fruit of camphor tree, *Pharmaceutical Journal*, No.6, (1991) 8.

- [2] N.X. Dung, P.V. Khiem, H.T. Chien, P.A. Leclercq, The essential oil of *Cinnamomum camphora* (L.) Sieb. var. *linaloolifera* from Vietnam, *J. Essent. Oil Res.*, 5 (1993) 451.
- [3] Nguyen Xuan Dung, Do Quang Huy, Pham Van Khiem, etc, *Pharmaceutical Journal*, No.4, (1988) 13.
- [4] Hoang Hai Hiên, Le Thi Mai Hoa, Tran Dinh Thang, Nguyen Xuan Dung, Some results of the study on Lauraceae Species from Vietnam, *Proceeding of The 9th International Congress on Ethnopharmacology*, Nanning, China, 22-26/8/2006.
- [5] Tran Dinh Thang, Do Ngoc Dai, Do Quang Huy, Nguyen Xuan Dung, A new natural source of Camphor from *Cinnamomum longepetiolatum* Costerm. apud Phamh. in Vietnam, *VNU Journal of Science, Nat., Sci., & Tech.*(Vietnam), 2008 (accepted).
- [6] Tran Dinh Thang, Do Ngoc Dai, Do Quang Huy, Nguyen Xuan Dung, Chemical composition of the leaf oil of *Cinnamomum ovatum* Allen from Vietnam, *Pharmaceutical Journal*, Vol.13, No.2, (2008) 94.
- [7] Tran Dinh Thang, Hoang Hai Hien, Trinh Xuan Thuy, Nguyen Xuan Dung, Volatile Constituents of the Leaf Oil of *Litsea euosma* J. J. Sm. from Vietnam, *Journal of Essential Oil Bearing Plants*, Vol.9, No.2 (2006) 122.
- [8] Nguyen Hoang Anh, Helmut Ripperger, Tran Van Sung, Guenter Adam, Neolignans and a sesquiterpene from *Caryodaphnosis tonkinensis*, *Phytochemistry*, 42 (4) (1996) 1167.
- [9] L. Zhu, D. Ding, B. M. Lawrence, The cinnamomum species in China: resources for the present and future, *Perfumer and Flavorist*, 19 (1994) 17.
- [10] R.P. Adams, *Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectrometry*, Allured Publishing Corp. Carol Stream, IL., 2001.
- [11] Yoshida, T., Muraki, S. and Komatsu, A.. Minor Constituents of Japanese Ho-Leaf Oil. *Agr. Biol. Chem.* 33 (1969) 343.
- [12] National Sun Yat-Sun University, *A systematic study on the genus Machilus of Taiwan* (Lauraceae), 1997.

## Thành phần hóa học của tinh dầu cây *Machilus Japonica* Siebold & Zucc. (Lauraceae) ở Việt Nam

Phạm Văn Khiên<sup>1</sup>, Đỗ Quang Huy<sup>2</sup>, Đỗ Thị Việt Hương<sup>2</sup>

<sup>1</sup>Tạp chí Dược học

<sup>2</sup>Trường Đại học Khoa học Tự nhiên, ĐHQGHN, 334 Nguyễn Trãi, Hà Nội, Việt Nam

Thành phần hóa học của cây *Machilus Japonica* Siebold & Zucc. (Lauraceae) ở Việt Nam đã được nghiên cứu. Tinh dầu thu được từ lá cây 3-4 năm tuổi bằng phương pháp cất lôi cuốn hơi nước. Thành phần hóa học của tinh dầu được xác định bằng phương pháp sắc ký khí và sắc ký khí khối phổ. Thành phần chính có mặt trong tinh dầu là  $\alpha$ -phellandrene (60,2%),  $\beta$ -phellandrene (10,5%), p-cymene (20,3%),  $\beta$ -myrcene (3,7%), terpineole (0,9%), trans- $\beta$ -ocimene (0,7%), 11-dodecenol (0,3),  $\beta$ -Caryophyllene (0,5%). Thành phần tinh dầu của loài cây lấy nghiên cứu chưa thấy có trong các công trình công bố trước đây ở Việt Nam. Kết quả nghiên cứu đã chỉ ra rằng đây là loài thuộc họ Lauraceae và có tên khoa học là *Machilus japonica* Siebold & Zucc.